## **Specimen Label**





### **Fungicide**

\*Trademark of Dow AgroSciences LLC

Active Ingredient

 Fenbuconazole: a-[2-(4-chlorophenyl)ethyl] 75.0%

 a-phenyl-1<u>H</u>-1,2,4-triazole-1-propanenitrile
 25.0%

 Inert Ingredients
 25.0%

 Total
 100.0%

U.S. Patent No. 4,366,165

EPA Reg. No. 62719-421

# Keep Out Of Reach Of Children CAUTION

### **Precautionary Statements**

#### **Hazards to Humans and Domestic Animals**

**CAUTION: Harmful If Swallowed Or Absorbed Through The Skin** 

Avoid contact with skin, eyes or clothing.

#### **Personal Protective Equipment (PPE)**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category **A** on an EPA chemical resistance category selection chart.

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber, or butyl rubber
- Chemical-resistant headgear for overhead exposure

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems enclosed cabs or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### **User Safety Recommendations**

#### Users should:

 Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

#### First Aid

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**Hot Line Number:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-992-5994 day or night, for emergency treatment information.

#### **Environmental Hazards**

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies elsewhere on this label. If terms are unacceptable, return at once unopened.

In case of emergency endangering health or the environment involving this product, call 1-800-992-5994. If you wish to obtain additional product information, visit our web site at www.dowagro.com.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

#### **Directions for use**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

#### **Agricultural Use Requirements:**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE)and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material such as polyvinyl chloride, nitrile rubber, or butyl rubber

#### Storage and Disposal:

**Storage:** Store in a cool, dry area above freezing. The water-soluble pouch may become brittle at storage temperatures below 32°F but the fungicide is not affected. Do not remove the water-soluble pouches from container except for immediate use.

**Pesticide Disposal:** Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environment Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

**Container Disposal**: Completely empty overwrap into application equipment then dispose of overwrap in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Steps to be Taken in Case Material is Released or Spilled: Keep dust to a minimum. Appropriate protective equipment must be worn when handling a spill of this material. Remove contaminated clothing promptly. Wash exposed skin areas with soap and water immediately after exposure. Contain spills immediately with inert materials (e.g. sand, earth). Transfer spilled material and solid diking material to separate suitable containers for recovery or disposal. Keep spills and cleaning runoff out of municipal sewers and open bodies of water. Refer to Material Safety Data Sheet.

#### General Information

Indar\* 75WSP fungicide is a protectant fungicide for the control of specific diseases mentioned on this label. Indar 75WSP fungicide should be used in a regularly scheduled preventive spray program. A wetting agent such as Latron B-1956\*, Latron CS-7\* or other non-polymer containing spray adjuvant approved for use with registered pesticide products should be added to spray solutions according to manufacturers' use instructions to achieve optimum disease control.

#### **Use Rate Determination**

Carefully read, understand and follow label use rates and restrictions. Under low disease conditions, minimum suggested use rates can be used, while maximum suggested label rates and shortened spray schedules are recommended for severe or threatening disease conditions. For proper application, determine the number of acres to be treated, the suggested label use rate and the gallonage to be applied per acre. Prepare only

the amount of spray solution required to treat the measured acreage. Careful calibration of spray equipment is recommended prior to use.

#### Mixing

Always place Indar 75WSP fungicide into solution prior to adding co-applied agricultural chemicals registered for use on specific crop or crops being treated. Add the required number of unopened pouches as determined by the dosage recommendations into the spray tank with agitation. Depending on the water temperature and the degree of agitation, the pouches should dissolve completely within approximately ten minutes from the time they are added to the water.

#### Handling

The enclosed pouches of Indar 75WSP fungicide are water-soluble. **Do not** allow pouches to become wet prior to adding to the spray tank. **Do not** handle the pouches with wet hands or wet gloves. Always reseal overwrap bag to protect remaining unused pouches. Do not remove water-soluble pouches from overwrap except to add directly to the spray tank.

#### **Application**

Do not make ground or aerial applications within 75 feet of streams, rivers, ponds, lakes or reservoirs.

**Ground:** Thorough coverage sprays generally result in optimum disease control. To achieve good coverage use proper spray pressure, gallonage per acre, nozzles, nozzle spacing and tractor speed. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration.

**Aerial:** For those crops where applications are not limited to ground equipment, apply in a minimum of 10 gallons of water per acre. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals. Flagmen and loaders should wash thoroughly before eating, drinking or smoking and at the end of each days spray operation.

## Aerial Spray Drift Management: AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

- 1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
- 2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed. The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory.

**Chemigation:** Do not apply this product through any type of irrigation system.

#### Compatibility

Indar 75WSP fungicide is compatible with most commonly used agriculture fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. When preparing tank mixes, user should consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use.

**Note:** Indar 75WSP fungicide is compatible with boron and spray oils; however, the water-soluble pouches must be completely dissolved before adding spray oils or products containing boron to spray-mixtures.

#### **Use Directions for Stonefruits**

Use one 2-ounce pouch of Indar 75WSP per acre in a minimum of 50 gallons by ground or 10 gallons by air. Indar 75WSP fungicide may be applied up to the day of harvest. Indar 75WSP is a protectant fungicide. Best disease control is achieved when a protectant application schedule is followed.

A wetting agent such as Latron B-1956, Latron CS-7 or other nonpolymer containing spray adjuvant approved for use in registered pesticides products should be added to spray solutions according to manufacturers' use instructions to achieve optimum disease control.

Crop	Disease	Use Recommendations
Apricots	Blossom Blight (Monilinia spp.)	Begin applications at early red bud stage before infection occurs.  If conditions are favorable for disease development, apply again at full bloom and at petal-fall.
	Fruit Brown Rot (Monilinia spp.)	Begin applications 2 to 3 weeks before harvest using a 7 to 10 day spray interval.
Cherries	Blossom Blight Fruit Brown Rot	Refer to apricots.
	Leaf Spot (Blumeriella spp.)	Follow blossom blight schedule and continue application at 10 to 14 day intervals. Additional foliar applications may be made after harvest.
Nectarines	Blossom Blight Fruit Brown Rot	Refer to apricots.
Peaches	Blossom Blight Fruit Brown Rot	Refer to apricots.
	Scab (Cladosporium spp.)	Begin applications at shuck split.  Make 2 to 3 subsequent thorough coverage applications at 10 to 14 day intervals.

#### **Restrictions (For All Crops):**

- Do not apply more than 1 lb (0.75 lb active) Indar 75WSP fungicide per acre per season.
- Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.

#### **Terms and Conditions of Use**

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

#### **Warranty Disclaimer**

Dow AgroSciences warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. Dow AgroSciences MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

#### **Inherent Risks of Use**

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of Dow AgroSciences or the seller. All such risks shall be assumed by buyer.

#### **Limitation of Remedies**

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Dow AgroSciences' election, one of the following:

- Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used

Dow AgroSciences shall not be liable for losses or damages resulting from handling or use of this product unless Dow AgroSciences is promptly notified of such loss or damage in writing. In no case shall Dow AgroSciences be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Dow AgroSciences or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

\*Trademark of Dow AgroSciences LLC

Dow AgroSciences LLC • Indianapolis, IN 46268 U.S.A.

Label Code: D02-845-003 Replaces Label: D02-845-002 LOES Number: 010-00341

EPA-accepted date: 03-26-2003

#### Revisions:

- Changes EPA-accepted 03-18-03:
   Revised First Aid statements in accordance with PR Notice 2001-1.
   Revised glove statement under PPE and in Agricultural Use Requirements box to read "Chemical-resistant gloves made of any waterproof material, such as polyvinyl chloride, nitrile rubber or lead to the base." butyl rubber."